

REPORT DOCUMENTATION PAGE

AD-A267 372



ORT DATE

3 REPORT TYPE AND DATES COVERED

ANNUAL 1 Jun 92 TO 31 May 93

4 TITLE AND SUBTITLE

(FY91 AASERT), RESEARCH TRAINING OF THE EFFECTS OF
TOXIC SUBSTANCES ON THE LUNGS

6 AUTHOR(S)

Dr Mark L. Witten

5 FUNDING NUMBERS

F49620-92-J-0325
61103D
3484
S4

7 PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Dept of Pediatric & Physiology
University of Arizona
1501 N. Campbell Avenue
Tucson, AZ 857248. PERFORMING ORGANIZATION
REPORT NUMBER

9 SPONSORING MONITORING AGENCY NAME(S) AND ADDRESS(ES)

AFOSR/NL
110 Duncan Avenue, Suite B115
Bolling AFB DC 20332-000110. SPONSORING MONITORING
AGENCY REPORT NUMBER

Dr Kozumbo

11. SUPPLEMENTARY NOTES

93-17267



12a. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release;
distribution unlimited

2b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

Allison M. Hays is the graduate student supported by the AASERT grant and Jason T. Figueroa was the high school student supported by the same grant. Jason was supported by the AASERT grant during the Summer of 1992. He worked on two research projects during this time period. The first project concerned the effects of JP-8 jet fuel exposure on substance P receptors, specifically the NK1 receptor, in rats' lungs. This project encountered technical difficulties in determining the sensitivity of the assay used in quantifying the amount of NK1 receptors. Consequently, we did not attempt to publish this data. Jason also worked on our acute smoke exposure project which is an established model of Acute Respiratory Distress Syndrome. His work with this project resulted in an abstract which he presented at the Experimental Biology '93 meeting in New Orleans. Jason was awarded a Flinn Foundation scholarship for this Fall to attend the Univ of Arizona and decided to terminate his association with the AASERT training grant. I have replace Jason with Brian Tollinger, a student who has just been accepted into the doctoral program in the College of Pharmacy. Brian is investigating the possibility of entering a combined Ph.D. in Toxicology- Pharm. D. program. Brian has worked in my laboratory for

14. SUBJECT TERMS

15. NUMBER OF PAGES

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT

(U)

18. SECURITY CLASSIFICATION
OF THIS PAGE

(U)


19. SECURITY CLASSIFICATION
OF ABSTRACT

(U)

20. LIMITATION OF ABSTRACT

(UL)

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

FIRST YEAR SUMMARY FOR AASERT GRANT
ENTITLED
RESEARCH TRAINING OF THE EFFECTS OF TOXIC SUBSTANCES
ON THE LUNGS

Mark L. Witten, Ph.D. Principal Investigator

Department of Pediatrics
Arizona Health Sciences Center
Tucson, Arizona

Accession No.	
NTIS Doc #	
DTIC No.	
U.S. GPO	
By	
Date	
A	
Dist	
A	

DTIC QUALITY INSPECTED 3

Submitted to-
Life and Environmental Sciences Directorate
U.S. Air Force Office of Scientific Research
Bolling Air Force Base, DC 20332-6448

25 JUN 1993

Over-all Progress of Grant

Allison M. Hays is the graduate student supported by the AASERT grant and Jason T. Figueroa was the high school student supported by the same grant. Jason was supported by the AASERT grant during the Summer of 1992. He worked on two research projects during this time period. The first project concerned the effects of JP-8 jet fuel exposure on substance P receptors, specifically the NK1 receptor, in rats' lungs. This project encountered technical difficulties in determining the sensitivity of the assay used in quantifying the amount of NK1 receptors. Consequently, we did not attempt to publish this data. Jason also worked on our acute smoke exposure project which is an established model of Acute Respiratory Distress Syndrome. His work with this project resulted in an abstract which he presented at the Experimental Biology '93 meeting in New Orleans. Jason was awarded a Flinn Foundation scholarship for this Fall to attend the University of Arizona and decided to terminate his association with the AASERT training grant. I have replaced Jason with Brian Tollinger, a student who has just been accepted into the doctoral program in the College of Pharmacy. Brian is investigating the possibility of entering a combined Ph.D. in Toxicology- Pharm. D. program. Brian has worked in my laboratory for the past year. His research project is to develop an assay for determining neutral endopeptidase concentrations in rat lung lavage and tissue samples after chronic exposure to JP-8 jet fuel using high pressure liquid chromatography.

Allison has worked on the JP-8 jet fuel inhalation toxicology project for the past two years under my direction and that of Dr. John K. Pfaff, a Navy physician who has conducted a post-doctoral Fellowship in my laboratory. Dr. Pfaff has been assigned duty at the Portsmouth Naval Hospital starting in July of 1993. Allison will assume Dr. Pfaff's duties on the JP-8 jet fuel inhalation toxicology project. Allison has also been heavily involved in our acute smoke exposure model of Acute Respiratory Distress Syndrome. This work resulted in an abstract that she presented at the Experimental Biology '93 meeting in New Orleans. Allison is doing well in her course work in preparation for entering the Doctoral program in the Department of Anatomy. However, she may transfer to the Department of Pharmacology/Toxicology doctoral program because she is concerned about future employment possibilities with a degree in Anatomy -vs- a degree in Toxicology. There is no doubt that a

Toxicology degree has many more employment possibilities than a degree in Anatomy at this point in time.

Plans for Year 2 of the Grant

Allison will take over the daily management of the Air Force JP-8 jet fuel project with Brian's assistance. Both Allison and Brian are intelligent students and I expect them to maintain their high level of performance in their coursework. Both will continue their individual research projects on the jet fuel project and I expect both of them to present their research at the Experimental Biology '94 meeting in Anaheim, California.

2938

PULMONARY NEUTROPHIL SEQUESTRATION AFTER INTESTINAL ISCHEMIA-REPERFUSION. M. Taha, B. Gerson, University of Chicago, Chicago, Illinois 60637.
Intestinal ischemia reperfusion is a known precipitant of acute lung injury. Neutrophils (PMN) and free radicals from PMN are critical to both processes. This study was designed to determine if the filtration of PMN from intestinal perfusate could attenuate lung PMN accumulations and injury. Four groups of rats (n=5 each) were subjected to 40 or 45 min of complete ischemia (superior mesenteric artery occlusion and collateral ligation). The SMA was then reperfused for 40 min with or without a leukocyte filter which removes 99% PMN. Sham (n=5) underwent identical preparation without ischemia or filter (ischemic time = 0). After 30 min of reperfusion, lung tissue was assessed for PMN accumulation (myeloperoxidase levels) and alveolar capillary leak (1.25 albumin). Filtering reperfusion PMN reduced lung PMN accumulation at 40 min after reperfusion in rats suffering both 40 and 45 min of ischemia. In contrast, there was no increase in lung microvascular permeability at the same post-reperfusion interval.

Lung MPO (ΔA/min/gm)

Ischemic Time

	7 min	40 min	45 min
No Filter	32.2 ± 3.5	71.7 ± 7.8*	136.5 ± 7.8*
Filter		40.6 ± 10.8	69.8 ± 11.9*

* p<0.05 from 0

/ p<0.05 from Filter

These data show that lung PMN accumulation occurs in the early period after intestinal reperfusion and predates functional evidence of lung injury. While a cause and effect relationship remains unproved, the known deleterious effects of PMN activation suggests that reperfusion filtration may have clinical application in decreasing distant organ dysfunction after intestinal ischemia.

Supported by NIH Surgical Scientist Training Grant

ENVIRONMENTAL PATHOLOGY AND AIR POLLUTANTS (1939-1942)

2939

MECHANISMS OF SMOKELESS TOBACCO-INDUCED INCREASE IN MICROVASCULAR PERMEABILITY IN VIVO. J. Rubinstein, Y. P. Gao, J. M. Conlon and J. K. Vishwanatha, Univ. of NE Medical Center and Creighton University, Omaha, NE 68198.

The purpose of this study was to determine whether smokeless tobacco (SE) increases vascular permeability in the hamster cheek pouch, and whether these effects are mediated by local generation and release of bradykinin (BK), coupled with a decrease in tissue angiotensin I-converting enzyme (ACE) activity, which cleaves BK. Using intravital microscopy, we found that SE extract induced a significant (p < 0.05) concentration-dependent increase in leaky site formation and clearance of fluorescein isothiocyanate dextran (m.w. = 70,000 daltons) in the cheek pouch. These effects were significantly attenuated by two selective bradykinin B₂ receptors antagonists, Hoe 140 and NPC 17647. Suffusion of SE extract was also associated with a significant increase in BK concentration in the suffusate, and with a significant decrease in cheek pouch ACE activity. Levels of ACE protein and mRNA in the cheek pouch were not altered during suffusion of SE extract. We conclude that SE extract increases microvascular permeability in the hamster cheek pouch by local generation and release of BK, and by decreasing tissue ACE activity leading to potentiation of BK-induced responses.

2941

DOES LEAD PLAY A ROLE IN THE PATHOGENESIS OF METABOLIC BONE DISEASE? A HYPOTHESIS. H. Spencer, V. Q. Sullivan and S. Somay, Metabolic Research, V. A. Hospital, Hines, IL 60141.

We reported previously that 89% of patients with Paget's disease of bone gave a history of occupational exposure to lead (J. Lab. Clin. Med., 120:798-800, 1992). Bone is the prime target organ of the deposition and long-term storage of lead. The question arose whether lead may also play a role in metabolic bone diseases other than Paget's disease. That this may apply to hyperparathyroidism was considered. Reason: others have shown that lead interferes with the utilization of vitamin D and induces vitamin D deficiency which would lead to decreased intestinal absorption of calcium, a low calcium status and subsequent parathyroid stimulation. The histories of 4 patients with proven hyperparathyroidism revealed that they were occupationally or environmentally exposed to lead for many years. A 6th patient, on whom no occupational history was obtained, may have been exposed to lead as he had both Paget's disease and hyperparathyroidism. We are extending our series at present. These preliminary observations should stimulate others to investigate whether lead is one of the factors which may play a role in the pathogenesis of the metabolic bone disease hyperparathyroidism.

2940

U75412E PRETREATMENT BEFORE ACUTE SMOKE EXPOSURE CAUSES A LARGE INCREASE IN LUNG PROSTACYCLIN CONCENTRATIONS. A.M. Hays, R.C. Lantz, M. Vermeulen, G. Chen, M.L. Wines, Steele Memorial Children's Research Center, University of Arizona, Tucson, AZ and Harvard Medical School, Boston, MA.

We utilized a rabbit model to analyze the effect of the lipoaroid, U75412E, on its ability to attenuate severe lung injury induced by acute smoke exposure. The acute smoke insult consisted of 60 tidal volume breaths of diesel fuel-polycarbonate plastic smoke administered in 8-9 min. There were four groups of rabbits: Three hour smoke-exposed rabbits (THSE, N=8), Three hour sham smoke-exposed rabbits (THSS, N=6), Short (3-4 min) smoke-exposed rabbits (SSE, N=5), and U75412E pretreated smoke-exposed (USE, N=7). The lungs were removed immediately after the experiment and bronchoalveolar lavage (BAL) was performed with normal sterile saline. BAL fluid was analyzed for 6-keto-PGF₁α, the stable metabolite of prostacyclin. The 6-keto-PGF₁α concentrations (pg/ml BAL fluid) were the following:

THSE	506 (32)
THSS	307 (57)
SSE	142 (25)
USE	1495 (107)*

* p < 0.05. The USE rabbits lived for six hours after the smoke insult before they were killed for cell culture studies. Perhaps, the increased lung prostacyclin production in the USE rabbits contributes to an attenuation of the smoke-induced lung injury by modulation of the airway bronchoconstrictive response to smoke. Supported by Upjohn.

2942

ADVERSE HEALTH EFFECTS ASSOCIATED WITH CHROMIUM EXPOSURE IN WORKERS EMPLOYED IN A CHROMATE INDUSTRY. D.M. Backyavathy and N.V. Nanda Kumar, SPQM, M.K.D. Pagala, P.G. Department of Zoology, Voorhees College, Madras University, Vellore 632 001 and Department of Zoology, Sri Venkateswara University, Tirupati 517 502, INDIA

Adverse health effects of Chromium in Industrial Workers exposed to occupational environment have been reported. Four hundred workers of different age groups employed in a chromate factory, showed occurrence of skin diseases like chrome-hold ulcer (28.75%), industrial dermatitis (18.75%), acid burns (9.5%), injuries (13.75%) and respiratory diseases like bronchitis (77.5%), acute pharyngitis (86.25%), fibrosis (2.5%), pleurisy (3%) and also nasal irritation and nasal perforation (18.25%) with high relative risk factor. The above adverse health effects are associated with occupational environment of the chromate factory. Total chromium content in Urine and Blood in workers ranged from 6.2-17.5 µg/L urine and 2.4-12.7 µg/100 G blood respectively.

2360

Acrolein and other aldehydes present in tobacco smoke induce activation/inactivation of protein kinase C - protection by thiol agents. **D. Jantzen, L. Gundimeda, Z. H. Chen, and R. Gopalakrishna**, Dept. Pharmacol. & Nutr., USC Sch. of Med., Los Angeles, CA 90033

Acrolein, formaldehyde, and acetaldehyde are toxic agents present in tobacco smoke and in polluted air. These aldehydes induce pathobiological effects in lungs including tumor promotion. Since protein kinase C (PKC) is a receptor for tumor promoters, we have determined whether these aldehydes could influence PKC. In the bronchial epithelial cells and NRK cells treated with acrolein (0.5 to 5 μ M) a rapid (min) 2- to 2.5-fold increase in PKC activity without any change in subcellular distribution was observed. However, prolonged treatment with acrolein resulted in an inactivation of PKC. These changes were also induced in purified PKC by direct exposure to acrolein. Both formaldehyde and acetaldehyde also induced similar cellular changes in PKC at much higher concentrations (>100 μ M). N-acetylcysteine, L-cysteine, and a cell-permeable analogue of glutathione all protected PKC from the aldehyde-induced modifications of PKC in intact cells. Thus, acrolein and other aldehydes induce pathobiological effects and tumor promotion in lungs, in part, by inducing activation/inactivation of PKC. Nonetheless, they differed from other tumor promoters (phorbol esters, oxidants) in that these aldehydes did not induce a cytosol to membrane translocation of PKC.

Supported by Grant RT388 TRDRP, University of California

2361

INCREASING SMOKE EXPOSURE PRODUCES ALVEOLAR EDEMA BY AIRWAYS NOT ALVEOLAR EDEMA WITHOUT LIPID PEROXIDATION. **J. P. Jantzen, D. Jantzen, L. Gundimeda, R. Deming, Beth Israel Hospital, 330 Brookline Avenue, Boston, MA 02215**

We determined the effect of a graded exposure to smoke exposure on lung physiology. Respective to alveolar edema induced by cigarette smoke (L.P. 10) sheep were given 12 breaths of cigarette smoke at a tidal volume of 1 ml, 10 ml, 20 ml, 40 ml, 80 ml, 160 ml, 320 ml, 640 ml, 1280 ml, 2560 ml, 5120 ml, 10240 ml, 20480 ml, 40960 ml, 81920 ml, 163840 ml, 327680 ml, 655360 ml, 1310720 ml, 2621440 ml, 5242880 ml, 10485760 ml, 20971520 ml, 41943040 ml, 83886080 ml, 167772160 ml, 335544320 ml, 671088640 ml, 1342177280 ml, 2684354560 ml, 5368709120 ml, 10737418240 ml, 21474836480 ml, 42949672960 ml, 85899345920 ml, 171798691840 ml, 343597383680 ml, 687194767360 ml, 1374389534720 ml, 2748779069440 ml, 5497558138880 ml, 10995116277760 ml, 21990232555520 ml, 43980465111040 ml, 87960930222080 ml, 175921860444160 ml, 351843720888320 ml, 703687441776640 ml, 1407374883553280 ml, 2814749767106560 ml, 5629499534213120 ml, 11258999068426240 ml, 22517998136852480 ml, 45035996273704960 ml, 90071992547409920 ml, 180143985094819840 ml, 360287970189639680 ml, 720575940379279360 ml, 1441151880758558720 ml, 2882303761517117440 ml, 5764607523034234880 ml, 11529215046068469760 ml, 23058430092136939520 ml, 46116860184273879040 ml, 92233720368547758080 ml, 184467440737095516160 ml, 368934881474191032320 ml, 737869762948382064640 ml, 1475739525896764129280 ml, 2951479051793528258560 ml, 5902958103587056517120 ml, 11805916207174113034240 ml, 23611832414348226068480 ml, 47223664828696452136960 ml, 94447329657392904273920 ml, 188894659314785808547840 ml, 377789318629571617095680 ml, 755578637259143234191360 ml, 1511157274518286468382720 ml, 3022314549036572936765440 ml, 6044629098073145873530880 ml, 12089258196146291747061760 ml, 24178516392292583494123520 ml, 48357032784585166988247040 ml, 96714065569170333976494080 ml, 193428131138340667952988160 ml, 386856262276681335905976320 ml, 773712524553362671811952640 ml, 1547425049106725343623905280 ml, 3094850098213450687247810560 ml, 6189700196426901374495621120 ml, 12379400392853802748991242240 ml, 24758800785707605497982484480 ml, 49517601571415210995964968960 ml, 99035203142830421991929937920 ml, 198070406285660843983859875840 ml, 396140812571321687967719751680 ml, 792281625142643375935439503360 ml, 1584563250285286751870879006720 ml, 3169126500570573503741758013440 ml, 6338253001141147007483516026880 ml, 12676506002282294014967032053760 ml, 25353012004564588029934064107520 ml, 50706024009129176059868128215040 ml, 101412048018258352119736256430080 ml, 202824096036516704239472512860160 ml, 405648192073033408478945025720320 ml, 811296384146066816957890051440640 ml, 1622592768292133633915780102881280 ml, 3245185536584267267831560205762560 ml, 6490371073168534535663120411525120 ml, 12980742146337069071326240823050240 ml, 25961484292674138142652481646100480 ml, 51922968585348276285304963292200960 ml, 103845937170696552570609926584401920 ml, 207691874341393105141219853168803840 ml, 415383748682786210282439706337607680 ml, 830767497365572420564879412675215360 ml, 1661534994731144841129758825350430720 ml, 3323069989462289682259517650700861440 ml, 6646139978924579364519035301401722880 ml, 13292279957849158729038070602803445760 ml, 26584559915698317458076141205606891520 ml, 53169119831396634916152282411213783040 ml, 106338239662793269832304564822427566080 ml, 212676479325586539664609129644855132160 ml, 425352958651173079329218259289710264320 ml, 850705917302346158658436518579420528640 ml, 1701411834604692317316873037158841057280 ml, 3402823669209384634633746074317682114560 ml, 6805647338418769269267492148635364229120 ml, 13611294676837538538534984297270728458240 ml, 27222589353675077077069968594541456916480 ml, 54445178707350154154139937189082913832960 ml, 108890357414700308308279874378165827665920 ml, 217780714829400616616559748756331655331840 ml, 435561429658801233233119497512663310663680 ml, 871122859317602466466238995025326621327360 ml, 1742245718635204932932477990050653242654720 ml, 3484491437270409865864955980101306485309440 ml, 6968982874540819731729911960202612970618880 ml, 13937965749081639463459823920405225941237760 ml, 27875931498163278926919647840810451882475520 ml, 55751862996326557853839295681620903764951040 ml, 111503725992653115707678591363241807529902080 ml, 223007451985306231415357182726483615059804160 ml, 446014903970612462830714365452967230119608320 ml, 892029807941224925661428730905934460239216640 ml, 1784059615882449851322857461811868920478433280 ml, 3568119231764899702645714923623737840956866560 ml, 7136238463529799405291429847247475681913733120 ml, 14272476927059598810582859694494951363827466240 ml, 28544953854119197621165719388989902727654932480 ml, 57089907708238395242331438777979805455309864960 ml, 114179815416476790484662877555959610910619729920 ml, 228359630832953580969325755111919221821239459840 ml, 456719261665907161938651510223838443642478919680 ml, 913438523331814323877303020447676887284957839360 ml, 1826877046663628647754606040895353774569915678720 ml, 3653754093327257295509212081790707549139831357440 ml, 7307508186654514591018424163581415098279662714880 ml, 14615016373309029182036848327162830196559325429760 ml, 29230032746618058364073696654325660393118650859520 ml, 58460065493236116728147393308651320786237301719040 ml, 116920130986472233456294786617302641572474603438080 ml, 233840261972944466912589573234605283144949206876160 ml, 467680523945888933825179146469210566289898413752320 ml, 935361047891777867650358292938421132579796827504640 ml, 1870722095783555735300716585876842265159593655009280 ml, 3741444191567111470601433171753684530319187310018560 ml, 7482888383134222941202866343507369060638374620037120 ml, 14965776766268445882405732687014738121276749240074240 ml, 29931553532536891764811465374029476242553498480148480 ml, 59863107065073783529622930748058952485106996960296960 ml, 119726214130147567059245861496117904970213993920593920 ml, 239452428260295134118491722992235809940427987841187840 ml, 478904856520590268236983445984471619880855975682375680 ml, 957809713041180536473966891968943239761711951364751360 ml, 1915619426082361072947933783937886479523423902729502720 ml, 3831238852164722145895867567875772959046847805459005440 ml, 7662477704329444291791735135751545918093695610918010880 ml, 15324955408658888583583470271503091836187391221836021760 ml, 3064991081731777716716694054300618367237478244367204320 ml, 6129982163463555433433388108601236734474956488734408640 ml, 12259964326927110866866776217202473468949912977468817280 ml, 24519928653854221733733552434404946937899825954937634560 ml, 49039857307708443467467104868809893875799651909875269120 ml, 98079714615416886934934209737619787751599303819750538240 ml, 196159429230833773869868419475239575503198607639501076480 ml, 392318858461667547739736838950479151006397215279002152960 ml, 784637716923335095479473677900958302012794430558004305920 ml, 1569275433846670190958947355801916604025588861116008611840 ml, 313855086769334038191789471160383320805117772222301723680 ml, 627710173538668076383578942320766641610235544444603447360 ml, 1255420347077336152767157884641533283220471088889206894720 ml, 25108406941546723055343157692830665664409421777784137440 ml, 50216813883093446110686315385661331328818843555568274880 ml, 100433627766186892221372630771322626657637687111136549760 ml, 200867255532373784442745261542645253315275374222273099520 ml, 40173451106474756888549052308529050663055074844454619840 ml, 80346902212949513777098104617058101326110149688909239680 ml, 160693804425899027554196209234116202652220299377818479360 ml, 321387608851798055108392418468232405304440598755636958720 ml, 642775217703596110216784836936464810608881197511273917440 ml, 1285550435407192220433569673872929621217762395022547834880 ml, 257110087081438444086713934774585924243552479004509566960 ml, 514220174162876888173427869549171848487104958009019133920 ml, 1028440348325753776346855739098343696974209916018038267840 ml, 2056880696651507552693711478196687393948419832036076535680 ml, 4113761393303015105387422956393374787896839664072153071360 ml, 8227522786606030210774845912786749575793679328144306142720 ml, 1645504557321206042154969182557349915158735865628861228440 ml, 3291009114642412084309938365114699830317471711257722456880 ml, 6582018229284824168619876730229399660634943422515444913760 ml, 1316403645856964833723975346045879932126988684503088982720 ml, 2632807291713929667447950692091759864253977369006177965440 ml, 5265614583427859334895901384183519728507954738012355930880 ml, 10531229166857718669791802768367039457015909476024711861760 ml, 21062458333715437339583605536734078914031818952049423723520 ml, 42124916667430874679167211073468157828063637904098847447040 ml, 84249833334861749358334422146936315656127275808197694894080 ml, 168499666669723498716668844293872631312254551616395399788160 ml, 336999333339446997433337688587745262624509103232790799576320 ml, 673998666678893994866675377175490525249018206465581599152640 ml, 1347997333357787989733350754350981050498036412931163195305280 ml, 2695994666715575979466701508701962100996072825862326390605440 ml, 5391989333431151958933403017403924201992145651724652781210880 ml, 10783978666862303917866806034807848403984291303449305562421760 ml, 2156795733372460783573361206961569680796858260689861112484320 ml, 4313591466744921567146722413923139361593716521379722224968640 ml, 862718293348984313429344482784627872318743304275944444977280 ml, 1725436586697968626858688965569255744637486608551888899544560 ml, 3450873173395937253717377931138511493274973217103777799089120 ml, 6901746346791874507434755862277022986549946434207555598178240 ml, 13803492693583749014869511724554045973099892868415111196356480 ml, 27606985387167498029739023449108091546199785736830222392712960 ml, 55213970774334996059478046898216183092399571473660444785425920 ml, 110427941548669992118956093796432366184799142947320889570851840 ml, 220855883097339984237912187592864732369598285894641779141703680 ml, 441711766194679968475824375185729464739196571789283558283407360 ml, 883423532389359936951648750371458929478393143578567116566814720 ml, 1766847064778719873903297500742917858956786287157134233133629440 ml, 3533694129557439747806595001485835717913572574314268466267258880 ml, 7067388259114879495613190002971671435827145148628853252534517760 ml, 14134776518229758991226380005943342871654290297257717055069035520 ml, 28269553036459517982452760011886685743308580594515434110138071040 ml, 56539106072919035964905520023773371486617161189030868220276142080 ml, 113078212145838071929811040047546742973234322378061736440552284160 ml, 226156424291676143859622080095093485946468644756123472881104568320 ml, 452312848583352287719244160190186971892937289512246945762209136640 ml, 904625697166704575438488320380373943785874579024493891524418273280 ml, 1809251394333409150876976640760747887571749158048987783048836546560 ml, 3618502788666818301753953281521495775143498316097975566097673093120 ml, 7237005577333636603507906563042991550286996632195951132195346186240 ml, 14474011154667273207015813126085983100573993264391902264390692372480 ml, 28948022309334546414031626252171966201147986528783804528781384744960 ml, 57896044618669092828063252504343932402295973057567609057562769489920 ml, 115792089237338185656126505008687864804591946115135218115125538979840 ml, 231584178474676371312253010017375729609183892230270436230251077959680 ml, 463168356949352742624506020034751459218367784460540872460502155919360 ml, 926336713898705485249012040069502918436735568921081744921004311838720 ml, 1852673427797410970498024080139005836873471378422163489842008623677440 ml, 3705346855594821940996048160278011673746942756844326979684017247354880 ml, 7410693711189643881992096320556023347493885513688653959360234944709760 ml, 14821387422379287763984192641112046694987771027377307918720469889419520 ml, 29642774844758575527968385282224093389975542054754615837440939778839040 ml, 59285549689517151055936770564448186779951084109509231674881879557678080 ml, 118571099379034302111873541128896373559902168219018463349763759115356160 ml, 237142198758068604223747082257792747119804336438036926699527518230712320 ml, 474284397516137208447494164515585494239608672876073853399055036461424640 ml, 94856879503227441689498832903117098847921734575214770

2121

ALTERATION IN ALVEOLAR MACROPHAGE FUNCTION FOLLOWING ACUTE SMOKE EXPOSURE. R.C. Lantz, G.J. Chen, A.M. Hays, M. Vermeulen and M. Witten. Depts. of Anatomy and Pediatrics, Univ. of Arizona Health Sciences Center, Tucson, AZ 85724 and Harvard Medical School, Boston, MA 02119.

New Zealand white rabbits were exposed to 60 tidal volumes of a synthetic smoke generated by the combustion of diesel fuel and polycarbonate plastic. Following exposure, animals were either sacrificed immediately (short term, N=5) or were maintained on a ventilator until they expired (2.95 ± 0.40 hrs) from the smoke exposure (long term, N=8). A third group was given a sham smoke exposure and sacrificed after 3 hrs (control, N=6). Pulmonary alveolar macrophages (PAM) were lavaged and cultured for up to 24 hrs. PAM were assessed for their ability to produce superoxide anion (O_2^-) and tumor necrosis factor alpha (TNF- α). Lavage fluid was also analyzed for the presence of TNF- α . PAM from long term animals had both increased basal and zymosan stimulated O_2^- production immediately after lavage. This elevated production persisted for at least 24 hrs in culture. O_2^- production in PAM from short term animals was not different than controls. Conversely, TNF- α production in PAM from long term animals was suppressed immediately following lavage and only returned to control levels after 24 hrs in culture. In contrast, TNF- α production from short term animals was significantly elevated over control values 2 hrs after lavage. While TNF- α levels in lavage fluid were elevated in long term animals, values were not significantly different from controls. We conclude that TNF- α production is stimulated in PAM immediately following smoke exposure while priming of PAM for O_2^- production is delayed. Sponsored by Upjohn Pharmaceutical Company and DoD Training Grant.

2123

Alteration of intracellular pH (pHi) and calcium ($[Ca^{2+}]_i$) in guinea pig alveolar macrophages during phagocytosis. Q.S. Qu, M. Ryan, and L.C. Chen. (SPON: C. Nadziejko) Dept. of Environmental Medicine, New York University Medical Center, Long Meadow Rd, Tuxedo, NY 10987.

The role of $[Ca^{2+}]_i$ and pHi in phagocytic function of guinea pig alveolar macrophages (AM ϕ) was investigated by measuring phagocytic index (PI, the percentage of viable AM ϕ that ingested at least one latex particle), phagocytic capacity (C, the percentage of actively phagocytizing AM ϕ that ingested four or more particles), pHi and $[Ca^{2+}]_i$ simultaneously in AM ϕ incubated with opsonized latex beads (3 μ m) at 0, 15, 30, 45, 60, and 90 minutes after incubation. Before exposed to latex beads AM ϕ were loaded with SNARF-1 and Fluo-3 together for pHi and $[Ca^{2+}]_i$ measurement. The resting $[Ca^{2+}]_i$ and pHi of AM ϕ was 64.85 ± 12.84 nM and 7.26 ± 0.08 (mean \pm SD; n=10), respectively. During incubation period, pHi decreased continuously with a concurrent increase in $[Ca^{2+}]_i$. The values of pHi and $[Ca^{2+}]_i$ measured at and after 60 minutes were significantly different from corresponding values measured when phagocytosis was first initiated ($p < 0.05$). The changes of pHi during phagocytosis correlate, to some extent, with changes of $[Ca^{2+}]_i$ ($r = 0.64$, $p = 0.0008$). Furthermore, phagocytic function of AM ϕ was closely associated with pHi and $[Ca^{2+}]_i$.

	PI		C	
	r	p	r	p
pHi	0.9123	0.0001	0.8436	0.0001
$[Ca^{2+}]_i$	0.6564	0.0042	0.7786	0.0015

Our data suggest that both pHi and $[Ca^{2+}]_i$ are important physiological parameters governing phagocytosis in AM ϕ of guinea pigs.

2125

PULMONARY SYSTEMIC HOST DEFENSE PHAGOCYTES IN THE RAT.

R. J. Frie, T. E. Hutson and G. D. Nisnam

Northeastern Ohio Universities College of Medicine, Rootstown, Ohio 44272

Bacteremia/endotoxemia stimulates an increase in the pulmonary host-defense capacity. The current study tested the hypothesis that microbial stimulation of the lung's systemic host-defense capacity is mediated by microvascular accumulation of activated mononuclear phagocytes. Sixteen male Sprague-Dawley rats (314 \pm 7 g) were anesthetized (10 mg Ketaset/100 g), equal volumes of the microbial product glucan (2 mg/100 g; n=9) or saline (n=7) infused via the dorsal penic vein and the animals permitted to recover. The rats were reanesthetized 48 hours later, intubated, and a catheter placed in the carotid artery. A test particulate, mononuclear blue B (9 mg/100 g), was infused and, 10 minutes later, the lungs inflated fixed with 2.5% glutaraldehyde (1 ml/100 g). Five random tissue samples were collected from each lung and ten high magnification (1000 \times magnification under oil) fields/sample were histologically evaluated (50 fields/lung) to define the number of mononuclear blue B containing phagocytes. The multi-lobular nucleus of the neutrophil was used to differentiate it from mononuclear phagocytes. The number of neutrophils per field were the same in saline (0.09 \pm 0.67) and glucan challenged (0.79 \pm 0.41) rats. The number of mononuclear phagocytes, however, significantly ($p < 0.05$) increased from 3.69 \pm 0.39 active cells/field in the saline challenged rats to 6.47 \pm 2.0 active phagocytes/field in the glucan challenged rats. The results are consistent with the hypothesis that mononuclear phagocytes are the primary systemic host-defense cells in the rat lung and that circulating microbial products stimulate an increased accumulation of these activated phagocytes in the lung microvasculature.

Support: American Heart Association, Ohio Affiliate - the studies plus Mr. Hutson's undergraduate Student Summer Research Fellowship; American Physiological Society - Mr. Frie's High School Teacher Summer Research Fellowship.

2122

LOSS OF VLA-4 EXPRESSION OCCURS WITH MONOCYTE DIFFERENTIATION INTO MACROPHAGES. E. McFadyen, L. Embree, J. M. Harlan, R. K. Albers. University of Washington Medical Center, Seattle, WA 98195.

We have shown that, unlike blood monocytes, alveolar macrophages do not express the β_1 integrin VLA-4. We hypothesized this loss of VLA-4 expression occurred as a result of monocyte differentiation into macrophages. The promyelocytic leukemia cell lines THP-1 and HL-60 were stimulated for 24 hours with Phorbol 12-Myristate 13-Acetate (PMA, 50 ng/ml), then cultured in RPMI + 10% FBS for up to 5 days. Cells were assayed daily for surface expression of VLA-4, VLA-5 and β_1 using flow cytometry and mRNA levels of VLA-4 via Northern analysis. Unstimulated cells were nonadherent and expressed VLA-4, VLA-5 and β_1 . Within 24 hr the cells became adherent. Flow cytometry showed decreased VLA-4 expression by 48 hr and its absence at 96 hr. VLA-4 mRNA decreased within 24 hr and was nearly undetectable by 72 hr. Surface expression of VLA-5 and β_1 were unchanged throughout all 5 days. Loss of VLA-4 may be important with regard to 1) marking monocyte differentiation into macrophages, 2) as a model for studying control of β_1 integrin expression and 3) mechanisms of monocyte recruitment from the vascular space.

2124

PULMONARY INTERSTITIAL MACROPHAGES (IM) AND DENDRITIC CELLS (DC) COOPERATE TO PROCESS PARTICULATE ANTIGEN. J. L. Gong, K. M. McCarthy, L. J. Tkachuk, E. E. Schneeberger. Massachusetts General Hospital, Boston, MA 02114.

Ia^b , FcR $^+$ interstitial lung DC, obtained at 95-98% purity from Lewis rats, do not endocytose PKH26 labeled heat killed *Listeria monocytogenes* (HKL), as visualized by fluorescence microscopy. In the total absence of macrophages, DC present this antigen poorly to HKL-immune T-cells (9,571 \pm 1,356 cpm) (mean \pm SE). Ia^b lung IM, obtained from the same rat lungs, avidly phagocytose PKH26 labeled HKL, but by themselves do not present HKL to HKL-immune T-cells (135 \pm 17 cpm). In an antigen presentation assay utilizing PKH26 labeled HKL, addition of IM (5 \times 10 4 /well) to lung DC (5 \times 10 4 /well), at a ratio of 1:10, results in a 300% increase in [3 H]Tdr uptake by HKL-immune T-cells (27,438 \pm 1,427 cpm). Addition of $>10\%$ IM to the total accessory cell population, however, produces a progressive inhibition of [3 H]Tdr uptake of HKL-immune T-cells. Supplementation with 10-30% (v/v) of 0.22 μ m filtered medium, derived from Ia^b IM (2.5 \times 10 4 /ml) that phagocytosed HKL for 16 h, results in a graded increase in [3 H]Tdr uptake of HKL-immune T-cells in the presence of DC. Supernatants derived from larger numbers of IM ($>10^4$ /ml), at the same IM:HKL ratio, are inhibitory when $>10\%$ (v/v) is added. It is concluded that small numbers of IM cooperate with DC to process and present particulate antigen; at higher numbers, however, IM inhibit antigen presentation by DC. Supported by NIH grant HL36781.

2126

CYTOKINE EXPRESSION BY MULTINUCLEATED ALVEOLAR MACROPHAGES INDUCED BY COLONY STIMULATING FACTORS. I. Lamare and H. Yang. Dept. of Pharm., Fac. Med., University of Ottawa, Ottawa, Ontario, Canada K1H 8M5.

Multinucleated giant alveolar macrophages (MGAM) are consistently found in lavage fluid and lung sections of animals with granulomatous inflammation. However, the mechanisms involved in MGAM formation and the precise role of these cells are unknown. In the present study, we investigated the *in vitro* effects of granulocyte-macrophage colony stimulating factor (GM-CSF) and macrophage colony stimulating factor (M-CSF) on rat alveolar macrophages (AM) cultured in Lab Tek chambers. Incubation with M-CSF (25-500 U/ml) caused within 3 days a significant increase in the number of MGAM with 3 or more nuclei that resemble those observed *in vivo*. M-CSF induced equally well morphologically different types of MGAM: a) MGAM with a round shape and 3-8 nuclei (Type 1), b) MGAM with irregular shape and 8-30 nuclei (Type 2). GM-CSF within the same dose range induced predominantly the formation of Type 2 MGAM. Interestingly, TNF expression was detected by immunocytochemistry in MGAM. Type 1 MGAM (80-90%) expressed high levels of cytoplasmic TNF whereas Type 2 (20-30%) expressed lower levels of TNF. These data indicate that M-CSF and GM-CSF may be involved in the formation of MGAM and that these cells are competent for TNF production (supported by MRC).